
No.



200200266

Anibersity of Idaho

DECEMS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT, VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANTS AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FERS AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN UCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY TION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (I) SHALL BE SOLD BY VARIETY NAME ONLY AS STIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER O TAT: 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT, COMMON

'Moreland'

In Testimonn Museon, I have hereunto set my hand and caused the seal of the Plant Mariety Protection Office to be affixed at the City of Washington, D.C. this twenty fourth day of June, year two thousand three.

Exhibit A, Origin and Breeding History of the Variety

Moreland was derived from a cross made at the University of Idaho, Aberdeen Research and Extension Center in 1986, designated A86327W with the pedigree 'Sonora 65'/II-60-155//'Heglar' (Cltr 17269) /3/'Warrior' (Cltr 13190)//'Kiowa'/PI 178383 /6/'Frocor'//'Frontana'/ 'Yaqui'/3/ 'Wanser' (CItr 13844) /4/ 'McCall' (CItr 13842) /5/Heglar /7/'WAID' (CItr 17806) /2*'Borah' (CItr 17267) //'Neeley' (CItr 17860). Frocor and II-60-155 are hard red spring wheat genotypes originating from Brazil and Minnesota, respectively. A86327W was advanced by the bulk method without intentional selection in the F2 generation. In the F3 generation, heads were selected from short plants and planted as F_{3,4} headrows in 1989. Heads from short plants were selected and advance by pedigree selection in 1990 and 1991 to form F_{6.7} headrows, which were harvested in 1992. From these headrows, the selection A86327W-3-2-2 was advanced to testing yield trials in southeastern Idaho in 1993. In 1996, A86327W-3-2-2 was designated IDO517 and entered into the Tri-State Irrigated Winter Wheat Nursery for two years. IDO517 was advanced into the Western Regional Winter Wheat Nursery for two years of testing (2000 and 2001). In 2001, IDO517 was evaluated in the Pacific Northwest Wheat Quality Council. IDO517 was also evaluated in the on-farm extension trials from 1999 to 2001 in Idaho and in 2001 in Oregon and Washington. In 2000, 200 head selections were planted at Aberdeen, ID and selected in summer 2001 for uniform plant type to form Moreland breeder seed. Seed from headrows that were true-to-type were harvested and planted at Aberdeen in 2001 to form foundation seed. IDO517 is uniform for plant type without obvious phenotypic variants and has remained uniform and stable for 11 generations during testing and seed increase, from 1992 to 2002.

Exhibit B, Novelty Statement

Moreland is most similar to the hard red winter wheat cultivar 'DW' developed by the University of Idaho. The two cultivars share a common parent, WAID/2*Borah//
Neeley. Moreland can be distinguished from DW on the basis of their reaction to dwarf bunt (causal organism *Tilletia controversa* Kühn in Rabenh). A summary of evaluations at Logan, UT are listed in Table 1.

Table 1. Reaction of hard red winter wheats to *Tilletia controversa* Kühn in Rabenh at Logan, UT as measured by percent of infected heads, averaged across two replications.

Year and trial	Moreland %	DW %	Average of adjacent susceptible check, Cheyenne
1997 Breeder trials	50	0	75
2000 Western regional nursery	10	0	17
2001 Western regional nursery	85	0	73
Average	47	0	55

Form Approved - OMB No. 0581-005:

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT C (Wheat)

OBJECTIVE DESCRIPTION OF VARIETY

	•	WHEAT (Tritic	cum spp.)	•
NAME OF APPLICANT(S)	,			FOR OFFICIAL USE ONLY
Universi	ty of Idaho	•		PVFO NUMBER
ADDRESS (Street and No. or RD N	lo., City, State, and Zip Code)			200200266
CALS Universi	ty of Idaho			VARIETY NAME
	ID 83844-2335			TEMPORARY OR EXPERIMENTAL DESIGNATION
a minimum of 100 plants. may be used to determine p	(c.g. 0 9 9 or 0 9) who	en number is either 99 or less or rmined from varieties entered in d:	r 9 or less respectively. n the same trial. Royal l	character of this variety in the boxes below. Data for quantitative plant characters should be based on Horticultural Society or any recognized color standard
1. KIND:				
1	1=Common	2=Durum	3=Club	4=Other (SPECIFY):
2. VERNALIZATIO	ON:			
2	1=Spring	2=Winter	3=Other (Sl	PECIFY) :
3. COLEOPTILE A	NTHOCYANIN:			3
1	1=Absent	2=Present		
4. JUVENILE PLAN	T GROWTH:		•	
2	1=Prostrate	2=Semi-erect	3=Erect	
5. PLANT COLOR	(boot stage):			
2	1 = Yellow-Green	2 = Green	3 = Blue-Gre	een
6. FLAG LEAF (boo	ot stage):		······································	· · · · · · · · · · · · · · · · · · ·
2	1 = Erect	2 = Recurved	2 may 3/16/2	1 = Not Twisted 2 = Twisted
7. EAR EMERGENO	CE:		•	1
0 4	Number of Days Earl	ier Than <u>Boundar</u>	у	*
	Number of Days Late	rThan <u>No rele</u>	vant earlie	er check *

S&T-470-6 (2-99) designed by the Plant Variety Protection Office with WordPerfect 6.0a Replaces LMGS-470-6 (6-82) which is obsolete

·	······································					
8. ANTHER CO	OLOR:				2002002)
1	1 = Yellow	2 = Pt	ırple		1500 1600 1500 1500 1600 1600 1600 1600	· ·
9. PLANT HEI	GHT (from soil to t	op of head, excludin	g awns):		MI	
1 8	cm Taller	Than Garland	<u>1</u>			*
1 0	cm Shorter	Than Boundar	<u>- y</u>		·	*
				* Relative to a DV/DO. As	pproved Commercial Variety Gro	our in the Come Take
10. STEM:				Relative to a 1 v r O-A	pproved Commercial Variety Gro	with the Same 1113
A. ANT	HOCYANIN	•		D. INTERNODE (SPI	ECIFY NUMBER)	
1	1= Absent 2=	Present		1 1= Hollow	2=Semi-solid	3=Solid
B. WAX	KY BLOOM		÷	E. PEDUNCLE		
1 1	l=Absent 2=	Present		1=Absent	2=Present	
С. НАП	RINESS (last interi	node of rachis)		9 cm Length		
1 1	(=Absent 2=)	Present				
11. HEAD (at M	Iaturity):					
A. DENS	SITY			C. CURVATURE		
	=Lax 2= = Dense	Middense		1 = Erect	2 = Inclined	3 = Recurved
B. SHAI	PE			D. AWNEDNESS		
		Strap Other (SPECIFY):		4 1 = Awnless 3 = Awnletted	2 = Apically Awnletted 4 = Awned	
12. GLUMES (a	t Maturity):				1 (1) (1) (1) (1) (1) (1) (1) (1) (1) (1	
A. COL				C. BEAK		
2 1	= White 2 =	- Tan		3 1 = Ob		
3	= Other (SPECIFY	7):		3 =Acu	minate	
B. SHOU	ULDER		*. *	D. LENGTH	·	
3 لــا	= Rounded 4 =	Oblique Square Apiculate			ort 2 = Medium 7mm) (ca. 8mm) ng (ca. 9mm)	
•						

*		A Chambart of a Cassacritica's
12.	GLUMES (at Maturity) Continued:	
	E. WIDTH	
	1 = Narrow (ca. 3mm) $3 = Wide (ca. 4mm)$	2 =

200200266

1	1 = Narrow (ca. 3mm)	2 = Medium (ca. 3.5mm
	3 = Wide (ca. 4mm)	

 _			

A. SHAPE

1 = Ovate

2 = Oval

C. BRUSH

1=Short

2=Medium

3=Long

1 = Not Collared

2 = Collared

B. CHEEK

1=Rounded

2=Angular

D. CREASE

1 = Width 60% or less of Kernel

2 = Width 80% or less of Kernel

3 = Width Nearly as Wide as Kernel

1 = Depth 20% or less of Kernel

2 = Depth 35% or less of Kernel

3 = Depth 50% or less of Kernel

G. PHENOL REACTION (see instructions):

E. Color

1=White. 2= Amber 4= OTHER (Specify)

3 = Red

3 = Elliptical

1 = Ivory

2 = Fawn

3 = Light Brown

4 = Dark Brown

5 = Black

F. TEXTURE

1=Hard

2=Soft

14. DISEASE:

(0=Not Tested;

1=Susceptible;

2=Resistant;

3=Intermediate;

4=Tolerant)

PLEASE INDICATE THE SPECIFIC RACE OR STRAIN TESTED

0	Stem Rust	(Puccinia graminis	f. sp. tritici)
---	-----------	--------------------	-----------------

Leaf Rust (Puccinia recondita f. sp. tritici)

Stripe Rust (Puccinia striiformis) 3

Loose Smut (Ustilago tritici) 2

Tan Spot (Pyrenophora tritici-repentis) 0

Flag Smut (Urocystis agropyri) 0

Halo Spot (Selenophoma donacis) 0

Common Bunt (Tilletia tritici or T. laevis) 1

Septoria nodorum (Glume Blotch) 3

Dwarf Bunt (Tilletia controversa)

Septoria avenae (Speckled Leaf Disease)

Karnal Bunt (Tilletia indica)

Septoria tritici (Speckled Leaf Blotch)

Powdery Mildew (Erysiphe graminis f. sp. tritici) 3

Scab (Fusarium spp.)

"Snow Molds" 0

14.	Disease (Continued)	(0=Not Tested;	1=Susceptible;	2=Resistant;	3=Interneli Je	0-00(Za6)6
		PLEASE IN	DICATE THE SP	ECIFIC RAC	E OR STRAIN TEST	TED
	2 "Black Point"	(Kernel Smudge)	C	Commo Bipolari	n Root Rot <i>(Fusariu</i> s spp.)	m, Cochliobolus and
	1 Barley Yellow	Dwarf Virus (BYDV) [Rhizocte	onia Root Rot <i>(Rhizo</i>	ctonia solani)
: .	O Soilborne Mos	aic Virus (SBMV)	4	Black C	haff <i>(Xanthomonas c</i>	ampestris pv. translucens)
	0 Wheat Yellow	(Spindle Streak) Mos	aic Virus	Bacteria syringae	il Leaf Blight <i>(Pseud</i>)	omonas syringae pv.
	1 Wheat Streak	Mosaic Virus (WSM	v) . [Other (SPECIFY)	
	Other (SPECI	FY)		Other (SPECIFY)	
· · · · · · · · · · · · · · · · · · ·	Other (SPECI	FY)		Other (S	SPECIFY)	•
	Other (SPECI	FY)		Other (S	SPECIFY)	
15. IN	SECT: (0=Not Test	ed; 1=Susceptible	; 2=Resistant;	3=Intermed	liate; 4=Tolerant)	
		PLEASE S	SPECIFY BIOTY	PE (where nee	eded)	
	1 Hessian Fly (M	ayetiola destructor)		Other (S	SPECIFY)	
	1 Stem Sawfly (C	Cephus spp.)		Other (S	SPECIFY)	
	1 Cereal Leaf Bee	tle <i>(Oulema melanop</i>	pa)	Other (S	SPECIFY)	
	7 Russian Aphid	(Diuraphis noxia		Other (S	SPECIFY)	
•	0 Greenbug (Sch	izaphis graminum)		Other (S	PECIFY)	
	0 Aphids			Other (S	PECIFY)	
16. AI	DITIONAL INFORMAT	TION ON ANY ITEM	I ABOVE, OR G	ENERAL CON	MMENTS	

Exhibit D, Additional Description of Variety

Moreland has an unpigmented coleoptile and semi-erect juvenile growth. Moreland has recurved, twisted flag leaf and an awned, erect, lax head, which is bronzechaffed at maturity. Moreland is 85 cm tall, similar to 'Brundage' soft white winter wheat, yet 10 cm shorter than 'Boundary' hard red winter wheat and 18 cm taller than 'Garland. Moreland is similar in heading date to 'Brundage', heading, on average in southern Idaho, 156 d after January 1. Moreland heads 4 d earlier than Boundary and 5 d earlier than Garland. Seed of Moreland is hard, red, ovate, and plump, with a kernel type similar to Neeley. Based on field evaluations in Washington and Idaho, Moreland has moderate adult plant resistance to stripe rust [caused by Puccinia striiformis (Westend.)]. Under severe stripe rust pressure at Pullman, WA, spring 2002, Moreland had a type 5 reaction occluding 30% of the flag leaf during grain fill. By comparison the resistant cultivar, Boundary had a type 5 reaction occluding 10% of the flag leaf, and the moderately susceptible cultivar Brundage had a type 8 reaction occluding 90% of the flag leaf. In five years of irrigated yield trials in southern Idaho, Moreland was more winter hardy than Boundary, Garland, and Stephens, with average spring stands of 99%, 96%, 96%, and 96%, respectively (LSD_{95%}=1%). Yet, Moreland has poorer snow mold (causal organisms Typhula spp.) tolerance than and greater susceptibility to dwarf bunt (Tilletia controversa Kühn in Rabenh) than Boundary.

Exhibit D, Additional Description of Variety (Continued).

Quantitative Data

- Table 2a. Agronomic Characteristics of Moreland hard red winter wheat in irrigated trials, 1997 to 2001.
- Table 2b. Agronomic Characteristics of Moreland hard red winter wheat in irrigated trials, 1994 to 2001.
- Table 2c. Results of Oregon-Idaho regional irrigated nursery, 2000 and 2001.
- Table 3. Agronomic performance of hard winter wheats in rain-fed trials, 2000 and 2001, six site-years.
- Table 4. Milling and baking data in irrigated research trials, southern Idaho, 1997 to 1999, total of 10 bakes, data by K. O'Brien.
- Table 5a to c. Stripe rust and infection type in winter cereals, 2000 and 2001, data of X. Chen, USDA-ARS.
- Table 6. Grain yield in 2001 extension trials in Oregon, data by R. Karow, OSU.
- Table 7. Washington State University hard winter wheat trials, extension testing program, 2001, data by J. Burns, WSU.
- Table 8. Paired comparisons of Moreland with hard red winter wheats in southern Idaho irrigated trials (Dist II, III, IV), 1999 to 2001, data by B. Brown and L. Robertson, UI.
- Table 9. Milling and baking data from rain-fed and irrigated extension trials in southern Idaho, 1999 to 2000, data by B. Brown, L. Robertson, and K. O'Brien.
- Table 10. Summary of the milling and baking performance of Moreland in the Western Regional Hard Nursery, 2000.

Table 2a. Agronomic Characteristics of Moreland hard red winter wheat in irrigated trials, 1997 to 2001

Aberdeen Hazelton

	Class	grain yield (bu/ac)	grain yield (bu/ac)	Test weight (lb/bu)	Spring stand (%)	Heading date (Julian)	Plant height (in)	Lodging (1=upright flat=9)
Moreland	HRW	145	129	60.1	99	156	34	1.0
Boundary	HRW	152	138	60.4	97	160	37	1.3
Garland	HRW	147	123	58.8	96	161	27	1.0
Brundage	SWW	157	135	60.4	96	157	33	1.0
Daws	sww	155	123	58.9	98	163	36	1.2
Stephens	sww	149	127	58.4	96	161	37	2.0
Std. Error		5.3	5.1	0.5	1	0.5	2	0.1

Table 2b. Agronomic Characteristics of Moreland hard red winter wheat in irrigated trials, 1994 to 2001

	Class	Aberdeen grain yield (bu/ac)	Hazelton grain yield (bu/ac)	Test weight (lb/bu)	Spring stand (%)	Heading date (Julian)	Plant height (in)	Lodging (1=upright flat=9)
Moreland	HRW	144	128	60.1	98	157	35	1.0
Garland	HRW	140	125	59.0	96	161	28	1.0
Stephens	sww	142	129	58.2	97	162	38	1.8
Daws	sww	147	124	59.0	97	163	37	1.4
Std error		3	3	0.3	5	0.5	1	0.2

Table 2c. Results of Oregon-Idaho regional irrigated nursery, 2000 and 2001

Hermiston, Parma, Hazelton, and Aberdeen

Cultivar	Grain Grain yield bu/ac	Test weight #/bu	Hazelton, ar Spring stand %	id Aberdeen Heading date julian	Height in
Moreland	137	61	97	148	32
Brundage	144	61	98	150	31
Stephens	135	59	98	154	35
Std. Error	ns	ns	ns	1	2

Table 3. Agronomic performance of hard winter wheats in rain-fed trials, 2000 and 2001, six site-years.

Cultivar	Grain yield bu/ac	Test weight #/bu	Spring stand %	Plant height in
Moreland	48	60.1	93	26
Bonneville	49	62.5	98	32
Boundary	52	59.7	96	27
Declo	48	60.7	94	25
Golden Spike	55	61.5	98	31
Promontory	48	63.2	97	29
Utah 100	53	60.4	96	32
Weston	51	63.2	95	34
Std error	3	1.5	3	2

Table 4. Milling and baking data in irrigated research trials, southern Idaho, 1997 to 1999, total of 10 bakes.

				Mixograph						
	Flour protein %	Milling yield %	Time to peak min	Height of peak cm	Mixing toler. Degrees	Water absorp.	Bake mix time min	Loaf volume ml	Interior texture 1 to 5	Exterior texture
Moreland	11.3	67.2	3.8	6.4	75.8	57.2	3.8	1009	4.1	16
Boundary	10.7	69.2	3.0	0.9	79.4	56.5	3.0	892	.	4.
Garland	11.1	64.2	2.2	6.5	70.9	57.4	2.1	935	1.2	2.
TSD 95%	9.0	1.0	0.3	0.4	2.4	SI	0.3	43	0.1	SU

TABLE 5a. STRIPE RUST PERCENT (%) AND INFECTION TYPE (T) ON CULTIVARS AND LINES IN THE WINTER REGIONAL DISEASE, 2001

NURSERY (WRDN, EXP01) AT SPILLMAN FARM (LOC01) AND WHITLOW FARM (LOC04) NEAR PULLMAN, WA AND MT VERNON, WA, (LOC5)

1					., , _ ~ .	<i></i> ,			
					STRIP	E RUS	ST T		
		LO	C01	LO	C04		LC	C05	
		Ī		4-	Jul	22-	Apr	21-	May
, , , , , , , , , , , , , , , , , , ,								STAG	E 4-5
CULTIVAR OR CR	PLOT	%	Τ	%	T	%	T	%	T
Bonneville	054	00	0	00	0	05	2	02	2
Boundary	163	00	0	02	5	05	8	10	2
Brundage	039	00	0	05	8	20	8	90	8
Moreland	Ave	00	8	01	8	10	8		
Gary	Ave	00	0	00	0	02	2		
Neeley	034	00	0	01	8	10	8		
Rod	055	00	0	02	8	10	8		2-5
Stephens	027	00	0	00	0	10	5		
WESTBRED 470	Ave	15	8	12	8			65	
	Bonneville Boundary Brundage Moreland Gary Neeley Rod	Bonneville 054 Boundary 163 Brundage 039 Moreland Ave Gary Ave Neeley 034 Rod 055 Stephens 027	CULTIVAR OR CRIPLOT %	LOC01 2-Jul STAGE 7 CULTIVAR OR CRIPLOT % T Bonneville 054 00 0 Boundary 163 00 0 Brundage 039 00 0 Moreland Ave 00 8 Gary Ave 00 0 Neeley 034 00 0 Rod 055 00 0 Stephens 027 00 0	LOC01 LOC01 LOC01 LOC01 LOC01 LOC01 LOC01 LOC01 STAGE 7 STAGE 7	STRIPI LOC01 LOC04 2-Jul 4-Jul STAGE 7 STAGE 6-7 S	LOC01 LOC04 2-Jul 4-Jul 22-STAGE 7 STAGE 6-7 STAGE 6	STRIPE RUST LOC01 LOC04 LOC04 LOC04 LOC04 LOC04 LOC04 STAGE 7 STAGE 6-7 STAGE 3 STAGE 3 STAGE 3 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 3 STAGE 3 STAGE 6-7 STAGE 6-7 STAGE 3 STAGE 3 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 6-7 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 6-7 STAGE 6-7 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 6-	STRIPE RUST LOC01 LOC04 LOC05 2-Jul 4-Jul 22-Apr 21-STAGE 7 STAGE 6-7 STAGE 3 STAGE 6-7 STAGE 6-7 STAGE 3 STAGE 6-7 STAG

TABLE 5b. STRIPE RUST PERCENT (%) AND INFECTION TYPE (T) ON CULTIVARS AND LINES IN THE WINTER REGIONAL DISEASE, 2000.

NURSERY (WRDN, EXP01) AT OBSERVATION HILL (LOC02) AND WHITLOW FARM (LOC04) NEAR PULLMAN, WA AND MT VERNON, WA, (LOC5)

							STRIP	E RU	ST			
				LO	C02		LO	C04		LOC	23	***
			7	lun	30-	Jun	28-	Jun	12-	-Apr	18-	May
		İ	STA	GE 7	STA	GE 8	STA	GE 6	STAC	SE 2-3	STA	GE 4
	CULTIVAR OR CR	PLOT	%	T	%	T	%	Τ	%	Τ	%	Τ
PI557014	Bonneville	054	00	0	00	0	00	0	05	2	05	2=5
PI14502B	Brundage	039	30	8	50	5=8	10	8	50	8	80	8
ID0517	Moreland	Ave	02	8	20	5=8	00	0	30	8	70	
IDO550	Gary	Ave	10	5	10	2=5	00	0	02	2	05	2=8
CI017860	Neeley	034	20	5	30	5	02	8	20			2=8
PI558510	Rod	055	30	8	30	5=8	00	0	30	8		2=8
CI017596	Stephens	027	00	0	05	2	00	0	10		_	2=8
PI599335	WESTBRED 470	Ave	70	8	90	8	20	8	30		90	

TABLE 5c. STRIPE RUST PERCENT (%) AND INFECTION TYPE (T) ON CULTIVARS AND LINES IN THE WINTER SOUZA DISEASE, 2001

NURSERY (EXP06) AT WHITLOW FARM (LOC04) NEAR PULLMAN, WA AND MT VERNON, WA, (LOC5) WHEN RECORDED AT THE

			Ç	TRIPE	RUS	T		
		LO	C04		LO	C05		
			Jul		Apr	22-	May	
		STAG	E 6-7	STA	GE 3	STAG	E 4-5	
ID NUMBER	PLOT	%	T	%	Т	%	Т	РМ
Boundary	800	00	0	05	2	10	5	
Brundage	010	05	8	30	8	80	8	
Garland	009	10	8	20	8	70	8	
Moreland	024	01	8	20	8	90	8	
Westbred 470	Ave	15	8	27	8	77	8	***-

Table 6. Grain yield in 2001 extension trials in Oregon.

	, ,			ogo			ı
			Grain yield	at individua	al locations		
0.145	0'	ļ					Average
Cultivar	Class	Corvallis	Hermiston	Madras	Ontario	Pendleton	grain
Moreland	HRW	163	116	135	125	103	128
Boundary	HRW	167	112	125	120	87	122
Delco	HRW	x	x	X	137	х	×
Bruehl	Club	152	99	110	56	81	100
Coda	Club	139	103	121	106	80	110
Edwin	Club	109	83	102	85	67	89
Rely	Club	148	81	121	109	95	111
Temple	Club	132	128	118	96	93	113
Hiller	Club	156	111	131	105	102	121
Rhode	Club	121	95	116	101	100	107
Connie	Durum	132	85	103	120	65	101
Brundage	SW	167	119	135	128	102	130
Foote	SW	162	101	117	96	98	115
Hubbard	SW	168	103	116	116	107	122
ID-B- 96	SW	142	112	137	129	97	123
Madsen	SW	164	108	120	123	85	120
Madsen/Stepl	n∈SW	174	114	128	123	96	127
Stephens	SW	154	128	113	138	100	127
Weatherford	SW	160	114	125	128	92	124
Yamhill	sw	155	113	107	102	83	112
Malcolm	SW	156	122	126	137	x	135
Rod	SW	169	136	121	122	96	129
Alzo	Trit	208	101	145	122	85	132
Bogo	Trit	178	88	132	143	86	125

Table 7. Washington State University hard winter wheat trials, extension testing program, 2001.

VARIETY NAME	Grain Yield	Test Weight	Grain Protein
Moreland	42.2	61.1	13.0
Hard White Common			
Gary	44.6	62.2	11.6
Hard Red Common			
BOUNDARY	44.7	61.0	12.4
BUCHANAN	44.1	61.7	11.9
COLUMBIA - 1	37.3	61.9	13.2
ESTICA	45.4	57.5	12.3
FINLEY	44.7	62.9	12.5
HATTON	41.2	63.4	12.5
RESIDENCE	51.9	60.0	12.0
SEMPER	50.0	59.7	12.1
SYMPHONY	44.5	60.7	13.0
WANSER	36.4	62.7	12.8
WESTON	45.0	63.0	13.2
Soft White Common			
ELTAN	48.2	61.5	11.8
NURSERY MEAN	43.3	61.9	12.4
CV %	14.4	0.9	4.5
LSD @ .10	2.8	0.3	0.3

Table 8. Paired comparisons of Moreland with hard red winter wheats in southern Idaho irrigated trials (Dist II, III, IV), 1999 to 2001.

0.10	- · ·	Grain yield	Test weight	Heading	Height	Lodging	Grain protein
Cultivar	Trials	bu/ac	#/bu	Julian	in	%	%
Moreland	12	126	61.9	152	33	0.6	11.7
Boundary	12	130	61.9	156	36	1.6	9.9
		ns	ns	**	***	ns	***
Moreland	18	128	62.3	152	34	0.3	11.2
Garland	18	121	61.6	155	28	0.3	11.2
	-	*	***	**	***	ns	ns
Moreland	10	123	61.5	150	34	0	11.2
Pillar	10	115	61.4	151	36	0.6	11.9
		***	ns	ns	*	ns	*
Moreland	15	132	63.1	151	35	0.3	11.2
Meridian	15	132	63.5	156	37	4.12	10.6
			ns	*	***	*	**

Table 10. Sum	Summary of the performance of Moreland in the Western Regional Hard Nursery, 20	erformano	e of Morela	nd in the We	stern Regi	onal Hard Ni	ursery, 2000		
Test Single K		Test	וויץ במסטומיי	Single	Kernel	Kernel	Grain		Break
		weight	Udy	kernel	weight	diameter	protein		flour
VARIETY	CLASS	#/bu	hardness	hardness	mg	mm	%		%
KHARKOF	HRW	62	70	59	31.7	2.2	14.4		38.4
WANSER	HRW	62.8	71	58.2	32.4	Ω	13.7		40.4
UT203032	HRW	62.7	78	59.3	36.4	2.5	12.8		41.4
NWCPOPE	HRW	රිය	68	57.4	အ	2.4	13.6		40.2
DECLO	HRW	62.4	82	70.3	30.1	2.2	13.7		32.6
Gary	WWH	62.9	78	62.2	33.6	2.4	12.7		38.7
Moreland	HRW	61	75	60.8	29.4	2.1	13.7	67.8	41.1
IDO562	HRW	64.8	87	59.2	36 3	2.5	12.9		42.2
MTW9441	WWH	62.7	75	63.9	27.5	2.1	13.6		41.9
MT9432	HRW	62.4	74	62.6	28.2	2.1	14.9		40.7
			Flour			Bake	Mix		
: !	I.	Milling	protein	RVA	Mix	absorp.	time	<u>vol</u>	Bread
VARIETY	FASH	score	%	units	type	%	min	≦	crumb
KHARKOT	0.38	80.1	12.7	158	3H	67.7	2.4	975	2
WANSER	0.37	84.2	12	167	4	67.8	3.3	1035	→
01203032	0.39	84.5	11.5	132	4	69.8	3.9	1065	_
NVCTCTE	0.4	81.5	11.9	173	2M	66.5	2.2	845	7
DECLO	0.43	75.9	11.5	155	웃	67.7	6.7	910	σı
cary	0.36	82.8	10.8	164	웃	65.5	5.5	920	4
Moreland	0.41	81.8	11.9	169	6H	66.5	5.6	930	ω
IDO562	0.37	86	11.2	157	5M	ල ල ග හ	ა 4	990	4
MTW9441	0.38	83.1	11.6	175	8	64.7	2.3	955	ພ.
W19432	0.36	84	12.7	184	5	68	3.9	955	ω

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in	n accordance with the Privacy Act
	1974 (5 U.S.C. 552a) and the Paperwo	rk Reduction Act (PRA) of 1995,
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to de certificate is to be issued (7 U.S.C. 2 until certificate is issued (7 U.S.C. 2426	421). Information is held confidentia
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
University of Idaho	OR EXPERIMENTAL NUMBER	
Idaho Agricultural Experiment Station	IDO517	Moreland
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
CALS	(208) 397-4181	(208)397-4311
University of Idaho	7. PVPO NUMBER	(200)337-4311
PO Box 442335 Moscow ID 83844-2335	7. PVPO NUMBER 2 0	0200266
		a m a m a m
8. Does the applicant own all rights to the variety? Mark an "X" in appropr	iate block. If no, please explain.	X YES NO
 Is the applicant (individual or company) a U.S. national or U.S. based co If no, give name of country 	ompany?	X YES NO
10. Is the applicant the original owner?	If no, please answer one of the fo	ollowing:
 a. If original rights to variety were owned by individual(s), is (are) the original 	iginal owner(s) a U.S. national(s)?	
☐ YES ☐ N	O If no, give name of country	\
b. If original rights to variety were owned by a company(ies), is(are) the	original owner(s) a U.S. based company	?
☐ YES ☐ N	O If no, give name of country	•
11. Additional explanation on ownership (if needed, use reverse for extra sp		
Moreland was derived from a single of the Idaho Agricultural Experiment Sta was performed under the direction of employee of the University of Idaho.	etion. Selection of t Professor Edward Sou Under the terms of f	he cultivar za, an
contract, the intellectual property of property of the University of Idaho.	created by faculty is	the
PLEASE NOTE:		
Plant variety protection can be afforded only to owners (not licensees) who meet on	e of the following criteria:	
1. If the rights to the variety are owned by the original breeder, that person must be which affords similar protection to nationals of the U.S. for the same genus and s	a U.S. national, national of a UPOV membe species.	er country, or national of a country
2. If the rights to the variety are owned by the company which employed the origin member country, or owned by nationals of a country which affords similar protection.	al breeder(s), the company must be U.S. base tion to nationals of the U.S. for the same go	ed, owned by nationals of a UPOV crus and species.
3. If the applicant is an owner who is not the original owner, both the original owner	r and the applicant must meet one of the abo	ove criteria.
The original breeder/owner may be the individual or company who directed final bre	eeding. See Section 41(a)(2) of the Plant Va	riety Protection Act for definition.

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to compete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require atternative means for communication of program information (braitle, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.